

10. BRYOCARPUM J. D. Hooker & Thomson, Hooker's J. Bot. Kew Gard. Misc. 9: 200. 1857.

长果报春属 *chang guo bao chun shu*

Herbs perennial. Leaves basal, petiolate. Flowers terminal, solitary, without bracts. Calyx 7-lobed to base; lobes lanceolate. Corolla yellow, funnellform-campanulate, exannulate, 7-lobed to middle; lobes linear, 2-lobed. Stamens 7, inserted toward apex of corolla tube; filaments very short; anthers oblong, apex acuminate. Ovary oblong, apex gradually narrowed into slender style. Stigma capitate. Capsule cylindric, many seeded, dehiscing by a lid at apex.

One species: Bhutan, China, Nepal, Sikkim.

1. *Bryocarpum himalaicum* J. D. Hooker & Thomson, Hooker's J. Bot. Kew Gard. Misc. 9: 200. 1857.

长果报春 *chang guo bao chun*

Rhizomes 1.5–5 cm, with oblong scales. Petiole 3.5–11 cm, usually longer than leaf blade; leaf blade ovate to ovate-oblong, (2.5–)3–7 × 2–3.5(–4) cm at anthesis, enlarged to 4.5–9 cm at fruiting, sparsely minutely dark glandular and short pubescent, base subrounded to cordulate, margin subentire to obscurely crenulate, apex obtuse to rounded. Scapes 1–3 per rosette, 10–20 cm, elongated to 35 cm in fruit. Calyx 7–12 mm, parted to base; lobes lanceolate, sparsely black glandular punctate. Corolla yellow, 1.5–2.5 cm, parted to middle; lobes linear, 2.5–4 mm wide, emarginate. Capsule 4–8 cm. Fl. May.

Pinus forests, mixed forests; 3000–4000 m. SE Xizang (M_{ཇོ}g Xian) [Bhutan, Nepal, Sikkim].

