164. STEREOCHILUS Lindley, J. Proc. Linn. Soc., Bot. 3: 38. 1858.

坚唇兰属 jian chun lan shu

Chen Xinqi (陈心启 Chen Sing-chi); Jeffrey J. Wood

Herbs, epiphytic, monopodial. Stems suberect, short or rather long, noded. Leaves several to many, distichous, flat, sometimes \pm conduplicate, leathery, jointed, sessile, often sheathing at base. Inflorescence 1–3, axillary, racemose, lax, often pendulous, many flowered; floral bracts very small, much shorter than pedicel and ovary. Flowers small, often opening widely; lateral sepals adnate to base of lip. Petals smaller than sepals; lip adnate to column base, immovable, saccately spurred at base, indistinctly 3-lobed; lateral lobes erect, small; mid-lobe larger; spur with longitudinal septum inside and often 1 or 2 calli on back wall. Column rather stout, without a foot; rostellum subulate-lanceolate, rather long; pollinia 4, in 2 pairs, equal, ellipsoid-oblong to ovoid, waxy, with short but distinct caudicles, attached by a long, slender, somewhat clavate stipe to a small, ovate viscidium.

Six described species: Bhutan, China, NE India, Myanmar, Thailand, Vietnam; two species in China, plus one undescribed species (see note below).

A third, as yet undescribed, apparently endemic species has been identified from S Yunnan, where it seems to be rather more common than either *Stereochilus brevirachis* or *S. dalatensis* (Perner, pers. comm.). It has the short inflorescence of *S. brevirachis* but has flowers with pink lip lateral lobes and a glabrous pedicel and ovary.

1a. Inflorescences shorter than leaves; pedicel and ovary publicent	1. S. brevirachis
1b. Inflorescences longer than leaves; inflorescence and ovary glabrous	2. S. dalatensis

1. Stereochilus brevirachis Christenson, Orchid Digest 62: 123. 1998.

短轴坚唇兰 duan zhou jian chun lan

Stems to 1.5 cm. Leaves 4–6, arcuate, linear, to 4.8×0.8 cm, dorsally carinate, apex obliquely minutely bilobed. Inflorescences laxly pendulous, somewhat zigzag, dark green, mottled purple, to 4.5 cm; floral bracts lanceolate, much shorter than ovary, acute. Flowers 2–5 per inflorescence; sepals, petals, and spur pale yellow to pale pink, lip mid-lobe magenta, lateral lobes yellow-orange, anther cap white with a central purple stripe; pedicel and ovary terete, to 12 mm, pubescent. Dorsal sepal elliptic, to 5×3 mm, concave, obtuse-rounded; lateral sepals obovate, to 5 × 4 mm, concave, obtuse-rounded. Petals oblong, to 6×2 mm, obtuse; lip 3-lobed; lateral lobes erect, transverse, abruptly narrowed and strongly incurved at apex; midlobe ovate, to 5×4 mm, obtuse, anterior callus a low transverse ridge at base, posterior callus a low keel from apex to spur to base of column surmounted by a large, reniform, bilaterally compressed callus; spur subparallel to ovary, ca. 4 mm, flattened on top and bottom, minutely bilobed at apex. Column ca. 3 mm; pollinarium to 4 mm. Fl. Jun.

Yunnan (without precise locality) [S Vietnam].

2. Stereochilus dalatensis (Guillaumin) Garay, Bot. Mus. Leafl. 23: 205. 1972.

坚唇兰 jian chun lan

Sarcanthus dalatensis Guillaumin, Bull. Mus. Natl. Hist. Nat., sér. 2, 27: 397. 1955.

Stems to 10 cm. Leaves to 12, distichous, dark green, often spotted purple, oblong-elliptic, ca. 5×0.5 cm, distinctly V-shaped in cross section, fleshy, apex rounded. Inflorescence axillary, borne from upper part of stem, 3–4 cm, laxly up to ca. 8-flowered; floral bracts ovate, minute, rounded. Flowers ca. 1 cm in diam.; sepals and petals white, lip light purple, hypochile yellow inside, anther cap deep purple; pedicel and ovary ca. 1 cm, glabrous. Sepals glabrous; dorsal sepal oblong-elliptic, ca. 5×3 mm, rounded; lateral sepals almost rhomboid, subspatulate, ca. 5×4 mm, 3-veined, rounded. Petals spatulate, ca. 5×4 mm, 1-veined, rounded; lip ligulate, ca. 5 mm; posterior callus papillose, sulcate in middle. Column ca. 4 mm; rostellum pendulous, long; anther cap narrowly triangular.

Yunnan (without precise locality) [Thailand, S Vietnam].

Flora of China 25: 463-464. 2009.