

落叶灌木；枝扁平。叶边缘有锯齿。花单一，稀2朵出自去年生枝的叶腋；花梗细长，与萼筒间无关节；花4出数，花冠4裂至基部，裂片反卷；花药背部无距。浆果4室。

已知2种，分布东亚、北美东部。我国1种，分布东部、中部至西南。

本组模式种：*V. erythrocarpum* (Pers.) Michx. (产北美东部)

91. 日本扁枝越桔

Vaccinium japonicum Miq. in Ann. Mus. Bot. Lugd. -Bat. **1**: 28. 1863; Maxim. in Bull. Acad. Sci. St. Petersb. **18**: 40. 1870; Franch. et Savat. Enum. Pl. Jap. **1**: 280. 1875; Rehd. in Journ. Arn. Arb. **15**: 288. 1934; Sleumer in Bot. Jahrb. **71**(4): 489. 1941; Chamberl. in Not. Roy. Bot. Gard. Edinb. **35**(2): 254. 1977; Ohwi, Fl. Jap. 1048. 1978.—*Oxycoccus japonicus* (Miq.) Makino in Bot. Mag. Tokyo **18**: 18. 1904.—*Oxycoccoides japonica* (Miq.) Nakai in Bot. Mag. Tokyo **31**: 247. 1917.—*Hugeria japonica* (Miq.) Nakai, Trees and Shrubs Japan ed. 2. 227. 1927; Maekawa in Bot. Mag. Tokyo **47**. fig. 14. 1933; Ohwi, Fl. Jap. 909. 1953; Makino, New Ill. Fl. Jap. 469. fig. 1875. 1979.

91a. 日本扁枝越桔(原变种)

var. *japonicum*

产日本。

91b. 扁枝越桔(中国高等植物图鉴)(变种) 扇木(江西)，山小檗(中国高等植物图鉴)，深红越桔、小叶梨状越桔(中国树木分类学) 图版39: 2—4

var. *sinicum* (Nakai) Rehd. in Journ. Arn. Arb. **5**: 56. 1924, in l. c. **15**: 288. 1934; Rehd. et Wils. in Journ. Arn. Arb. **8**: 185. 1927; Sleumer in Bot. Jahrb. **71**(4): 491. 1941; Chamberl. in Not. Roy. Bot. Gard. Edinb. **35**(2): 255. 1977.—*Oxycoccoides japonica* (Miq.) Nakai var. *sinica* Nakai, Trees and Shrubs Japan ed. 1. 168. 1922 et in Bot. Mag. Tokyo **38**: 40. 1924.—*Agapetes vaccinioidea* Lévl. in Bull. Acad. Géogr. Bot. **12**: 251. 1903, Fl. Kouy-Tchéou 147. 1914.—*Vaccinium siccum* Lévl. et Van. in Fedde, Repert. Sp. Nov. **9**: 447. 1911; id. Fl. Kouy-Tchéou 154. 1914.—*Hugeria sinica* (Nakai) Maekawa in Bot. Mag. Tokyo **47**: 615. fig. 16. 1933.—*H. japonica* (Miq.) Nakai var. *sinica* (Nakai) Hand.-Mazz. Symb. Sin. **7**(4): 799. 1936.—*H. vaccinioidea* (Lévl.) Hara in Journ. Jap. Bot. **21**: 146. 1947; 中国高等植物图鉴 **3**: 190. 图 4334. 1974.—*Vaccinium japonicum* auct. non Miq.: Forbes et Hemsl. in Journ. Linn. Soc. Bot. **26**: 16. 1889; Rehd. et Wils. in Sargent, Pl. Wils. **1**: 562. 1913; 陈嵘, 中国树木分类学 969. 1953.

落叶灌木，高0.4—2米；茎直立，多分枝，枝条扁平，绿色，无毛，有时有沟棱。叶散