

243):84. 1920. — Subject. *Pubescentes* Schneid. in Fedde, Rep. Sp. Nov. 9:80. 1910, nom. subnud., Ill. Handb. Laubh. 2:775. 1911. — Subser. *Pubescentes*(Schneid.) Rehd. in McKelvey, Lilac 11.1928.

单叶,全缘,叶片下面多少被毛。花序常由侧芽抽生,基部常无叶;花直径约6毫米;花药紫色、紫黑色或黄色。果多数具皮孔。

约4种,3亚种,2变种,1变型,我国均产,其中1种系栽培。

8. 巧玲花(中国高等植物图鉴) 小叶丁香(河南),雀舌花,毛丁香

Syringa pubescens Turcz. in Bull. Soc. Nat. Moscou 13:73. 1840; Hance in Journ. Bot. n. s. 4:133. 1875; Franch. in Bull. Soc. Philom. Paris ser. 7, 9:125. 1885; Bretschneid. Hist. Europ. Bot. Discov. China 351. 1057. 1898; Schneid. in Fedde, Rep. Sp. Nov. 9:80. 1910 et Ill. Handb. Laubh. 2:776, f. 486 v—y, 487 d—f. 1911; Bean. Trees & Shrubs Brit. Isles 2:571. 1914; Rehd. in Bailey, Stand. Cycl. Hort. 6:3300. 1917 et Bibl. Cult. Trees & Shrubs 566. 1949 et Man. Cult. Trees & Shrubs ed. 6, 781. 1954; Lingelsh. in Engl. Pflanzenr. 72(IV-243):84, f. 4. 1920, p.p. excl. syn. *S. tsinlingsana* Schneid. & *S. dielsiana* Schneid; McKelvey, Lilac 159, pl. 80—85. 1928; 陈嵘, 中国树木分类学 1051, 图 934. 1937; M. Noda, Fl. N.-E. Prov. China 1018. 1971; 中国高等植物图鉴 3: 350, 图 4654. 1974, excl. specim. gansu.; M. C. Chang & X. L. Chen in Investigat. Stud. Nat. 10:34. 1990. — *S. villosa* Vahl var. *β. ovalifolia* DC. Prodr. 8:283. 1844. — *S. villosa* var. *pubescens* Anonymous in Gard. Chron. ser. 3, 38:123, f. 43. 1905. — *S. pubescens* var. *typica* f. *pilosa* Schneid. in Fedde, Rep. Sp. Nov. 9:80. 1910 et Ill. Handb. Laubh. 2:776. 1911. — *S. wulingensis* Skv. & Wang in Ill. Fl. Lign. Pl. N.-E. China 478. 1955, Addenda 566. 1955. — *S. villosa* auct. non Vahl 1804:Decne. in Nouv. Arch. Mus. Hist. Nat. Paris ser. 2, 2:41. 1879; Franch. Pl. David. 1:204. 1884; J.D. Hook. in Curtis's Bot. Mag. 115:t. 7064. 1889; Hemsl. in Journ. Linn. Soc. Bot. 26:83. 1889; Dipp. Handb. Laubh. 1:116, f. 66. 1889.

8a. 巧玲花(原亚种) 图版 18:1—3

subsp. ***pubescens***

灌木,高1—4米;树皮灰褐色。小枝带四棱形,无毛,疏生皮孔。叶片卵形、椭圆状卵形、菱状卵形或卵圆形。长1.5—8厘米,宽1—5厘米,先端锐尖至渐尖或钝,基部宽楔形至圆形,叶缘具睫毛,上面深绿色,无毛,稀有疏被短柔毛,下面淡绿色,被短柔毛、柔毛至无毛,常沿叶脉或叶脉基部密被或疏被柔毛,或为须状柔毛;叶柄长0.5—2厘米,细弱,无毛或被柔毛。圆锥花序直立,通常由侧芽抽生,稀顶生,长5—16厘米,宽3—5厘米;花序轴与花梗、花萼略带紫红色,无毛,稀有略被柔毛或短柔毛;花序轴明显四棱形;花梗短;花